arXiv:math/0703075v1 [math.DG] 3 Mar 2007

GAUSSIAN CURVATURE AT THE HYPERELLIPTIC WEIERSTRASS POINTS.

ABEL CASTORENA

Abstract. Let C be a compact Riemann surface of genus $g > 0$. The principal polarization of the Jacobian $J(C)$ of C induces a metric on C of non-positive curvature, called the Theta metric. Let K be the Gaussian curvature of the Theta metric. We show that the hyperelliptic Weierstrass points are non-degenerated critical points of K of Morse index $+2$. We give an example in genus two where we show that the curvature K is a Morse function.

0.- Introduction and statement of results

Let C be a compact Riemann surface of genus $g > 0$. Consider the first homology group $H_1(C,\mathbb{Z}) \simeq \mathbb{Z}^{2g}$ and the complex g-dimensional vector space of holomorphic differentials $H^0(C,\Omega)$ and let $H^0(C,\Omega)^\vee$ be the dual space. The map $\jmath : H_1(C,\mathbb{Z}) \to H^0(C,\Omega)^\vee$, $\jmath(\gamma) = \int_\gamma$ is injective and the image generates a lattice in $H^0(C,\Omega)^\vee$. The Jacobian of C, $J(C) := H^0(C,\Omega)^\vee / H_1(C,\mathbb{Z})$ is a complex torus of dimension g. Take basis $\{\lambda_1,\ldots,\lambda_g,\beta_1,\ldots,\beta_g\}$ of $H_1(C,\mathbb{Z})$ and $\{\omega_1,\ldots,\omega_g\}$ of $H^0(C,\Omega)$, the $2g \times g$ period matrix of C with respect to this choice is $(\int_{\lambda_i}\omega_j, \int_{\beta_i}\omega_j), i,j = 1,...,g$.

Fixing a point $p_0 \in C$, we have in coordinates the Abel-Jacobi map $\mu_{p_0} : C \to J(C)$, $\mu_{p_0}(p) = (\int_{p_0}^{p}\omega_1,\ldots,\int_{p_0}^{p}\omega_g)$. Abel's theorem says that the Abel-Jacobi map is an embedding. The principal polarization of $J(C)$ is determined by the intersection form Q on $H_1(C,\mathbb{Z})$. When the basis of homology is symplectic, that is, $Q(\lambda_i,\lambda_j) = 0 = Q(\beta_i,\beta_j)$, $Q(\lambda_i,\beta_j) = \delta_{ij}$, $i,j = 1,\ldots,g$, there exists a unique basis $\{\omega_1,\ldots,\omega_g\}$ of $H^0(C,\Omega)$(called normalized basis) such that the period matrix of C is given by (I,Z) where I is the $g \times g$ identity matrix. From the Riemann bilinear relations([1], p.231-232) we have that Z is symmetric and

---

2000 *Mathematics Subject Classification.* 53A55

The author is partially supported with grant PAPIIT-IN104906-2, Universidad Nacional Autonoma de Mexico(UNAM), and grant 48668-F(CONACyT) Mexico.

that the hermitian form $(,)$ on $H^0(C,\Omega)$ defined by $(\eta,\theta) = i\int_C \eta\wedge\overline{\theta}$ satisfies $\frac{1}{2}(\omega_i,\omega_j) = \text{Im }\int_{\beta_i}\omega_j > 0$, that is, Im Z is positive definite. Let $(a_{ij}) = (\text{Im } Z)^{-1}$ be. In complex coordinates $(z_1,\dots,z_g)$ on $\mathbb{C}^g$ the Kähler form invariant under translations $\omega_0 := \sum_{i,j=1}^{g} a_{ij}dz_i\wedge d\overline{z}_j$ represents the principal polarization $[\omega_0]\in H^2(J(C),\mathbb{Z})$ of $J(C)$, that is, $[\omega_0]$ is the first Chern class of the line bundle defined by the Theta divisor of $J(C)$([1], p.333). The restriction of ω_0 to C give us a metric of non-positive curvature. We call this metric the Theta metric (other authors call this metric the Bergmann metric).

Let $p\in C$ and $z: U\subset C\to\mathbb{C}$ an holomorphic local coordinate around p. Let $f_i: U\to\mathbb{C}$ be a local expression of ω_i in the coordinate z, where $\{\omega_1,\dots,\omega_g\}$ is a basis of $H^0(C,\Omega)$ as above. We denote $f_i(z) := f_i\circ z^{-1}: z(U)\to\mathbb{C}$ and $\mathbf{f}(z) = (f_1(z),\dots,f_g(z))$. In this coordinate the Theta metric is given by $\rho^2(z)dz\wedge d\overline{z}$, where $\rho^2(z) :=< \mathbf{f}(z),\mathbf{f}(z) >= \mathbf{f}(z)\cdot(\text{Im } Z)^{-1}\cdot\overline{\mathbf{f}(z)^t} = (\sum\limits_{i,j=1}^{g} a_{ij}f_i(z)\overline{f_j}(z))$ is the conformality factor of the metric. The Gaussian curvature K is computed by the formula $K = \frac{-2}{\rho^2}\partial\overline{\partial}(\log(\rho^2))$ ([1], p. 77). Since $\frac{\partial}{\partial z}\rho^2(z) =< \mathbf{f}'(z),\mathbf{f}(z) >$, $\frac{\partial}{\partial\overline{z}}\rho^2(z) =< \mathbf{f}(z),\mathbf{f}'(z) >$, where $\mathbf{f}'(z) = (f_1'(z),\dots,f_g'(z))$, by the formula of the curvature we obtain:

$$K(z) = \frac{-2}{\rho^6(z)}(< \mathbf{f}(z),\mathbf{f}(z) >< \mathbf{f}'(z),\mathbf{f}'(z) > -| < \mathbf{f}'(z),\mathbf{f}(z) > |^2). \tag{1}$$

By Schwartz inequality we have that $K(p) = 0$ if and only if $\mathbf{f}(z)$ and $\mathbf{f}'(z)$ are linearly dependent. This happen if and only if the differential $d\phi$ of the canonical map $\phi: C\to\mathbb{P}(H^0(C,\Omega)^*) = \mathbb{P}^{g-1}$, $\phi(z) = [f_1(z):\cdots:f_g(z)]$ is zero in p. But $d\phi(p) = 0$ if and only if C is hyperelliptic and p is one of the $2g+2$ Weierstrass points. So in the hyperelliptic case the hyperelliptic Weierstrass points are global maxima of K.(See also [3]).

In this paper we prove:

Theorem. Let C be a hyperelliptic Riemann surface of genus $g > 0$. The Weierstrass points of C are non-degenerated critical points of Morse index $+2$ of the Gaussian curvature function K of the Theta metric.

In section two we show in lemma 2.1 and lemma 2.2 a criteria to find local minima for K. We give an example in genus two where we compute all critical points of the Curvature function K and we apply the theorem and lemmas 2.1, 2.2 to show that in this example K is a Morse function.

1. The Hessian of the Curvature function at the Weierstrass points.

Remark 1. Let C be a compact Riemann surface and $J(C)$ the Jacobian of C. Let $p_0, p_1 \in C, p_0 \neq p_1$ two points of C. Consider the Abel-Jacobi maps, $\mu_{p_0} : C \to J(C)$, $\mu_{p_1} : C \to J(C)$. Since $J(C)$ has structure of abelian group, for each $y \in J(C)$ we have the translation $\tau_y : J(C) \to J(C)$, $\tau_y(x) = x + y$. For $\mu_{p_1}(p_0) \in J(C)$ consider the translation $\tau_{\mu_{p_1}(p_0)}$. We have a commutative diagram:

$$\begin{tikzcd}
C \arrow[r, "\mu_{p_1}"] \arrow[dr, "\mu_{p_0}"'] & J(C) \\
& J(C) \arrow[u, "\tau_{\mu_{p_1}(p_0)}"']
\end{tikzcd}$$

that is, $\mu_{p_1} = \tau_{\mu_{p_1}(p_0)} \circ \mu_{p_0}$. Since ω_0 is invariant under translations we have that $\mu_{p_1}^* \omega_0 = \mu_{p_0}^* \tau_{\mu_{p_1}(p_0)}^* \omega_0 = \mu_{p_0}^* \omega_0$. Thus we have that Theta metric doesn't depend of the base in the Abel-Jacobi map.

The following lemma is useful to find critical points for K.

Lemma 1.1.- Let C be a compact Riemann surface of genus g and let $f : C \to C$ be an automorphism such that $f \neq$ identity. If $p \in C$ is a fixed point of f, then p is a critical point of the curvature function K of the Theta metric.

Proof. Let $f : C \to C$ be an automorphism, $f \neq$ identity. Let f_*, f^* be the induced isomorphisms in $H_1(C, \mathbb{Z})$ and $H^0(C, \Omega)$ respectively. Let $(f^*)^\vee : H^0(C, \Omega)^\vee \to H^0(C, \Omega)^\vee$ the dual map of f^*, where $H^0(C, \Omega)^\vee$ is the dual vector space to $H^0(C, \Omega)$. The following diagram commutes:

$$\begin{tikzcd}
H_1(C, \mathbb{Z}) \arrow[r, "f_*"] \arrow[d, "\jmath"'] & H_1(C, \mathbb{Z}) \arrow[d, "\jmath"'] \\
H^0(C, \Omega)^\vee \arrow[r, "(f^*)^\vee"] & H^0(C, \Omega)^\vee
\end{tikzcd}$$

where $\jmath$ is given by integration. This diagram induces an automorphism ϕ in the quotient $J(C) = H^0(C, \Omega)^\vee / H_1(C, \mathbb{Z})$ such that the following diagram commutes

$$\begin{tikzcd} C \arrow[r, "f"] \arrow[d, "\mu_p"'] & C \arrow[d, "\mu_{f(p)}"'] \\ J(C) \arrow[r, "\phi"] & J(C) \end{tikzcd} \tag{2}$$

For $\eta \in H^0(C,\Omega)$, $\; f^*(\overline{\eta}) = \overline{f^*(\eta)}$, then for $\theta_1, \theta_2 \in H^0(C,\Omega)$ we have:

$$(f^*\theta_1, f^*\theta_2) = \int\limits_C f^*\theta_1 \wedge \overline{f^*\theta_2} = \int\limits_{f_*(C)} (\theta_1 \wedge \overline{\theta_2}) = \int\limits_C (\theta_1 \wedge \overline{\theta_2}) = (\theta_1, \theta_2) \tag{3}$$

Then f^* is a isometry on $H^0(C,\Omega)$. Consider the dual hermitian inner product of the Theta metric, $(\cdot,\cdot)_1$, defined on $H^0(C,\Omega)^\vee$. Let $D(\phi)$ be the derivate of ϕ. The dual map of $D(\phi)$ is the dual map of f^*, hence $D(\phi)$ is an isometry with respect to $(\cdot,\cdot)_1$. Hence we deduce that ϕ leaves invariant the principal polarization ω_0. From (3) we have

$$\mu_p^*(\cdot,\cdot)_1 = \mu_p^*((\phi)^*(\cdot,\cdot)_1) = f^*\mu_{f(p)}^*((\cdot,\cdot)_1) = f^*\mu_{(p)}^*(\cdot,\cdot)_1.$$

Then f is an isometry with respect to the Theta metric. Hence f leaves invariant the curvature function K of the Theta metric, that is, $K(x) = K(f(x))$ for all $x \in C$. By the chain rule $K'(x) = K'(f(x))f'(x)$. If p is a fixed point of f then $K'(p) = K'(p)f'(p)$. But $f'(p) \neq 1$(apply Schwartz lemma to the lifting of f to universal covering spaces) and hence p is a critical point of K. ■

Example 1.- We recall that a hyperelliptic Riemann surface C of genus $g > 0$, is the compactification of the complex affine plane curve

$$C_0 := \{(x,y) \in \mathbb{C}^2 | y^2 = f(x),\; f(x) = (x-a_1)\cdots(x-a_{2g+2})\} \tag{$\bigstar$}$$

where $a_1, \ldots, a_{2g+2} \in \mathbb{C}$ and $a_i \neq a_j, \forall i \neq j$. The fixed points of the involution $\iota : C \to C$, $\iota(x,y) = (x,-y)$ are the $2g+2$ hyperelliptic Weierstrass points of C. So we obtain again that the Weierstrass hyperelliptic points are critical points of K. Note that the quotient $C/ < \iota >$ induces a double covering $\pi : C \to \mathbb{CP}^1$, $\pi(x,y) = x$ ramified at the Weierstrass points.■

Example 2.- Let C be the hyperelliptic Riemann surface given by the equation $y^2 = z^{2g+2} - 1$, and $\pi : C \to \mathbb{P}^1, \pi(z,y) = z$ the ramified double cover on $\mathbb{P}^1$. Consider the automorphism $h : C \to C$, $h(z,y) = (e^{\frac{2\pi i}{2g+2}}z, y)$. Note that

$(0,i), (0,-i)$ are fixed points of h, so they are critical points of K. Consider the automorphism $f : C \to C,\ f(z,y) = (\frac{1}{z}, \frac{iy}{z^{g+1}})$ which leaves invariant the curvature K of the Theta metric on C. f descends to $\mathbb{CP}^1$ and induces the automorphism $T : \mathbb{CP}^1 \to \mathbb{CP}^1, T(z) = \frac{1}{z}$. The following diagram commutes

$$\begin{tikzcd}
C \arrow[r, "f"] \arrow[d, "\pi"'] & C \arrow[d, "\pi"'] \\
\mathbb{CP}^1 \arrow[r, "T"] & \mathbb{CP}^1
\end{tikzcd}$$

Since $\pi : C \to \mathbb{P}^1$ is a double cover ramified at the Weierstrass points, we can take an open subset $U \subset \mathbb{CP}^1$ not containing the roots of $z^{2g+2} - 1 = 0$ such that $\pi^{-1}(U) = U_1 \cup U_2$ is a disjoint union where U_1, U_2 are open subsets of C and for $j = 1, 2$, $\pi|_{U_j} : U_j \to U$ is a local biholomorphism. We have that the open set $V := T(U)$ does not contain the roots of $z^{2g+2} - 1 = 0$, also $\pi^{-1}(V) = V_1 \cup V_2$ is a disjoint union and $\pi|_{V_j} : V_j \to V$ is local biholomorphism. Since $\pi|_{U_1} : U_1 \to U$ and $\pi|_{V_1} : V_1 \to V$ are local biholomorphism, we can restrict the Theta metric to the open set U_1, so we have that $T|_U = \pi|_{V_1} \circ f \circ \pi_{U_1}^{-1} : U \to V$ is an isometry with respect to the Theta metric on $\mathbb{P}^1$. Since f leaves invariant the curvature K, then T leaves invariant the curvature K on $\mathbb{P}^1$, that is, K descends to $\mathbb{P}^1$ and $K(z) = K(\frac{1}{z})$. Thus we obtain that the points lying over $0 \in \mathbb{P}^1$ and $\infty \in \mathbb{P}^1$ are critical points of K. ■

Now consider a hyperelliptic Riemann surface defined as in (★) in example 1. The Weierstrass point are the points $p_k = (a_k, 0)$, for $1 \leq k \leq 2g+2$. Note that $f'(a_k) \neq 0$, by implicit function theorem there exist a small neighbourhood U of p_k, such that y is a local coordinate. Let $g_k(x) = \prod_{i=1, i\neq k}^{2g+2}(x - a_i)$ be, and $z : C_0 \to \mathbb{C},\ z(x,y) = \frac{y}{\sqrt{g_k(x)}}$, in a neighbourhood of p_k. With this notation we have the following lemma:

Lemma 1.2.- z is a local coordinate in $p_k = (a_k, 0)$. In this coordinate the local expression of $(x - a_k)^{j-1}\frac{dx}{y}$ is $\alpha(z)z^{2(j-1)}dz$ for $j = 1, \ldots, g$, where $\alpha(z)$ is holomorphic and $\alpha(p_k) \neq 0$.

Proof. Since $dy = \frac{f'(x)dx}{2y} \neq 0$ in a neighbourhood of p_k, $g_k(a_k) \neq 0$ and $y(a_k) = 0$ then we have that $dz(p_k) = (\frac{\sqrt{g_k(x)}dy - yd\sqrt{g_k(x)}}{g_k(x)})(p_k) = (\frac{dy}{\sqrt{g_k(x)}})(p_k) \neq 0$, then there exists an open neighbourhood $\mathcal{V}$ of p_k where z is local coordinate. Note that $\frac{dx}{y} = \frac{2dy}{f'(x)}$, then for $j = 1, \ldots, g$ we have

$\frac{(x-a_k)^{j-1}dx}{y} = \frac{2(x-a_k)^{j-1}dy}{f'(x)} = z^{2(j-1)}(\frac{2dy}{dz}dz)\frac{1}{f'(x)} = \alpha(z)z^{2(j-1)}dz$

where $\alpha(z) = \frac{2y'(z)}{f'(x)}$ is a holomorphic function suh that $\alpha(p_k) \neq 0$. ■

Using translations we have that for $y \in J(C)$ we can identify the tangent space $T_y(J(C))$ with the tangent space at the origin $0 \in J(C)$, $T_0(J(C)) \simeq \mathbb{C}^g$. We use the coordinate $(z, \mathcal{V})$ of lemma 1.2 and we fix $q \in \mathcal{V}$. Consider the Abel-Jacobi map giving locally by

$$\mu : \mathcal{V} \subset C \to J(C)\ , \mu(z) = (\int_q^z \alpha(z), \int_q^z \alpha(z)z^2, \ldots, \int_q^z \alpha(z)z^{2(g-1)}).$$

The derivate of $\mu(z)$ is $\mu'(z) = \alpha(z)F(z)$, where $F(z) = (1, z^2, z^4, \ldots, z^{2(g-1)})$. Let $<,>$ any hermitian inner product on $T_0(J(C)) \simeq \mathbb{C}^g$ represented by an $g \times g$ hermitian symmetric matrix (h_{ij}) positive definite. In the coordinate $(z, \mathcal{V})$ the pullback $(\mu_q)^*(<,>)$ is a Kahler metric on C given by $\rho^2(z)dz \wedge d\overline{z}$, $\rho^2(z) = |\alpha(z)|^2\beta(z)$ where $\beta(z) =< F(z), F(z) >$. Let $G(z) = \beta(z)\frac{\partial^2\beta(z)}{\partial z \partial \overline{z}} - \frac{\partial\beta(z)}{\partial z} \cdot \frac{\partial\beta(z)}{\partial \overline{z}}$ be. Since $\alpha(z)$ is holomorphic, $\frac{\partial^2}{\partial\overline{z}\partial z}\log(\alpha(z)\overline{\alpha(z)}) = 0$. We have in this case that the curvature function on C with respect to this metric is given by $K(z) = \frac{-2G(z)}{\rho^2(z)}$.

Lemma 1.3. We have the following identities

$$\begin{aligned}
\frac{\partial G}{\partial z} &=< F', F >< F', F' > + < F, F >< F'', F' > - < F', F' >< F', F > \\
&\quad - < F, F' >< F'', F > . \\
\frac{\partial^2 G}{\partial z^2} &=< F'', F >< F', F' > + < F', F >< F'', F' > + < F', F >< F'', F' > \\
&\quad + < F, F >< F''', F' > - < F'', F' >< F', F > - < F', F' >< F'', F > \\
&\quad - < F', F' >< F'', F > - < F, F' >< F''', F > . \\
\frac{\partial^2 G}{\partial \overline{z} \partial z} &=< F', F >< F', F'' > + < F, F' >< F'', F' > + < F, F >< F'', F'' > \\
&\quad - < F', F'' >< F', F > - < F, F'' >< F'', F > - < F, F' >< F'', F' > .
\end{aligned}$$

Proof. It is clear that we have that $\frac{\partial\beta(z)}{\partial z} =< F'(z), F(z) >$, $\frac{\partial\beta(z)}{\partial\overline{z}} =< F(z), F'(z) >$, and

$$G(z) =< F(z), F(z) >< F'(z), F'(z) > - < F(z), F'(z) >< F'(z), F(z) > .$$

Now use the product rule for derivates. ■

Proof of the Theorem. We have

$$\begin{aligned}
&< F(p_k), F(p_k) >= h_{11},\\
&< F''(p_k), F''(p_k) >= 4h_{22}, \qquad (4)\\
&< F''(p_k), F(p_k) >= 2h_{12}.
\end{aligned}$$

From lemma 1.3 and (4), and the fact that $F'(p_k) = F'''(p_k) = (0, \ldots, 0)$ we obtain that

$$\frac{\partial^2 G}{\partial \overline{z} \partial z}(p_k) = 4(h_{11}h_{22} - h_{12}h_{21}), \quad \frac{\partial G}{\partial z}(p_k) = \frac{\partial^2 G}{\partial z^2}(p_k) = 0. \qquad (5)$$

We havel also,

$$\begin{aligned}
\frac{\partial K}{\partial z} &= \frac{-2}{\rho^2(z)} \cdot \frac{\partial G}{\partial z} + G \cdot \frac{\partial}{\partial z}(\frac{-2}{\rho^2(z)}).\\
\frac{\partial^2 K}{\partial z^2} &= \frac{-2}{\rho^2(z)} \cdot \frac{\partial^2 G}{\partial z^2} + 2\frac{\partial G}{\partial z} \cdot \frac{\partial}{\partial z}(\frac{-2}{\rho^2(z)}) + G \cdot \frac{\partial^2}{\partial z^2}(\frac{-2}{\rho^2(z)}).\\
\frac{\partial^2 K}{\partial \overline{z} \partial z} &= (\frac{-2}{\rho^2(z)}) \cdot \frac{\partial^2 G}{\partial \overline{z} \partial z} + \frac{\partial G}{\partial z} \cdot \frac{\partial}{\partial \overline{z}}(\frac{-2}{\rho^2(z)}) \qquad (6)\\
&\quad + G \cdot \frac{\partial^2}{\partial \overline{z} \partial z}(\frac{-2}{\rho^2(z)}) + (\frac{-2}{\rho^2(z)}) \cdot \frac{\partial G}{\partial \overline{z}}.
\end{aligned}$$

From lemma 1.3 and (4),(5), (6) and the fact that $G(p_k) = 0$ we have

$$\frac{\partial^2 K}{\partial z^2}(p_k) = 0 = \frac{\partial^2 K}{\partial \overline{z}^2}(p_k), \quad \frac{\partial^2 K}{\partial \overline{z} \partial z}(p_k) = \frac{-8(h_{11}h_{22} - h_{12}h_{21})}{\rho^2(p_k)}, \qquad (7)$$

note that $r := \frac{-8(h_{11}h_{22} - h_{12}h_{21})}{\rho^2(p_k)} < 0$. We have the matrix $\begin{pmatrix} \frac{\partial^2 K}{\partial z^2} & \frac{\partial^2 K}{\partial z \partial \overline{z}} \\ \frac{\partial^2 K}{\partial \overline{z} \partial z} & \frac{\partial^2 K}{\partial \overline{z}^2} \end{pmatrix}(p_k) = \begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix}$. Let $z = u + iv$ be. In real coordinates we have the following identities:

$$\begin{aligned}
\frac{\partial}{\partial z} &= \frac{1}{2}(\frac{\partial}{\partial u} - i\frac{\partial}{\partial v}), \frac{\partial}{\partial \overline{z}} = \frac{1}{2}(\frac{\partial}{\partial u} + i\frac{\partial}{\partial v})),\\
\frac{\partial^2}{\partial^2 u^2} &= 2\frac{\partial^2}{\partial z \partial \overline{z}} + (\frac{\partial^2}{\partial z^2} + \frac{\partial^2}{\partial \overline{z}^2}),\\
\frac{\partial^2}{\partial^2 v^2} &= 2\frac{\partial^2}{\partial z \partial \overline{z}} - (\frac{\partial^2}{\partial z^2} + \frac{\partial^2}{\partial \overline{z}^2}), \qquad (8)\\
\frac{\partial^2}{\partial u \partial v} &= \frac{1}{i}(\frac{\partial^2}{\partial \overline{z}^2} - \frac{\partial^2}{\partial z^2}),
\end{aligned}$$

Consider the matrix $\begin{pmatrix} \frac{\partial^2 K}{\partial u^2} & \frac{\partial^2 K}{\partial u \partial v} \\ \frac{\partial^2 K}{\partial v \partial u} & \frac{\partial^2 K}{\partial v^2} \end{pmatrix} := \begin{pmatrix} a & b \\ b & c \end{pmatrix}$. From (8) we have

$$\begin{pmatrix} \frac{\partial^2 K}{\partial z^2} & \frac{\partial^2 K}{\partial z \partial \overline{z}} \\ \frac{\partial^2 K}{\partial \overline{z} \partial z} & \frac{\partial^2 K}{\partial \overline{z}^2} \end{pmatrix} (p_k) = \begin{pmatrix} 0 & r \\ r & 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{4}(a-c-2ib) & \frac{1}{4}(a+c) \\ \frac{1}{4}(a+c) & \frac{1}{4}(a-c+2ib) \end{pmatrix}$$

The equations $\frac{1}{4}(a-c-2ib) = 0$, $\frac{1}{4}(a+c) = r$, $\frac{1}{4}(a-c+2ib) = 0$, implies that the real Hessian of K in p_k is Hess $K(p_k) = \begin{pmatrix} r & 0 \\ 0 & r \end{pmatrix}$. Hence for $1 \leq k \leq 2g+2$, the Weierstrass point $p_k = (a_k, 0)$ is a non-degenerated critical point of K of Morse index $+2$. ■

2.- An example in genus two.

Let C be the hyperelliptic Riemann surface of genus two defined by the equation $y^2 = (z-a_1)\cdots(z-a_6)$ with $a_i \neq a_j$. Take a simplectic homology basis $\{\lambda_1, \lambda_2, \beta_1, \beta_2\}$ in $H_1(C, \mathbb{Z})$. Consider the basis $\frac{dz}{y}, \frac{zdz}{y}$ of $H^0(C, \Omega)$. The period matrix with respect to this choice of basis is

$$\begin{pmatrix} \int_{\lambda_1} \frac{dz}{y} & \int_{\lambda_2} \frac{dz}{y} & \int_{\beta_1} \frac{dz}{y} & \int_{\beta_2} \frac{dz}{y} \\ \int_{\lambda_1} \frac{zdz}{y} & \int_{\lambda_2} \frac{zdz}{y} & \int_{\beta_1} \frac{zdz}{y} & \int_{\beta_2} \frac{zdz}{y} \end{pmatrix} := \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{14} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \alpha_{24} \end{pmatrix}$$

Since $\gamma := \alpha_{11}\alpha_{22} - \alpha_{12}\alpha_{21} \neq 0$([1], 231-232), we consider a normalized basis of differentials given for the change of basis:

$$\begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} := \frac{1}{\gamma} \begin{pmatrix} \alpha_{22} & -\alpha_{12} \\ -\alpha_{21} & \alpha_{11} \end{pmatrix} \cdot \begin{pmatrix} \frac{dz}{y} \\ \frac{zdz}{y} \end{pmatrix}$$

The period matrix of C with respect this new choice of differentials is (I, Z), where $Z = Z^t$ and Im $Z > 0$. Denote by $P := \frac{1}{\gamma} \begin{pmatrix} \alpha_{22} & -\alpha_{12} \\ -\alpha_{21} & \alpha_{11} \end{pmatrix}$. Let $\omega = (\omega_1, \omega_2)$ and $\eta = (\frac{dz}{y}, \frac{zdz}{y})$ be, then $\omega = \eta \cdot P^t$. Let $B := \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = P^t \cdot (\text{Im } Z)^{-1} \cdot \overline{P}$ be. The Theta metric on C is giving by $\rho^2(z)(dz \wedge d\overline{z}) = \omega \cdot (\text{Im } Z)^{-1} \cdot \overline{\omega^t} = \eta \cdot B \cdot \overline{\eta^t} = \frac{g(z)}{|y^2|} dz \wedge d\overline{z}$, where $g(z) = b_{11} + b_{21}z + b_{12}\overline{z} + b_{22}|z|^2$. Note that $b_{12} = \overline{b_{21}}$ and $\det B > 0$. With this notation we have the following lemma.

Lemma 2.1.- Let $(z_0, y_0) \in C$ be such that $f(z_0) \neq 0$, and $f'(z_0) = 0$. Then (z_0, y_0) is a critical point of K if and only if Re $b_{12} = 0$.

Proof. Let $(z_0, y_0) \in C$ such that $f(z_0) \neq 0$, and $f'(z_0) = 0$. We take a small open neighbourhood U_{z_0} of z_0 as coordinate of (z_0, y_0), we have a local biholomorphism $s : U_{z_0} \to \mathbb{C}$, $s(z) = z - z_0$. We consider in this coordinate the basis $\frac{ds}{y}, \frac{sds}{y}$. In this coordinate the conformality factor $\rho^2(z)$ of the Theta metric is given by $\frac{g(z)}{|y^2|}$ where $g(z) = b_{11} + b_{21}(z - z_0) + b_{12}\overline{(z - z_0)} + b_{22}|z - z_0|^2$. Since $f(z)$ is holomorphic, $\partial\overline{\partial}\log(f\overline{f}) = 0$, so $\partial\overline{\partial}\log(\frac{1}{|y^2|}) = \frac{1}{2}\partial\overline{\partial}\log((\frac{1}{|y^4|})) = \frac{1}{2}\partial\overline{\partial}\log((\frac{1}{f\overline{f}})) = 0$. The curvature function $K : U_{z_0} \to \mathbb{R}$ is given by:

$$K(z) = -2\frac{|y^2|}{g(z)} \cdot \partial\overline{\partial}\log(\frac{g(z)}{|y^2|}) = \frac{-1}{8}\partial\overline{\partial}\log(\frac{g(z)}{|y^2|}) \cdot \frac{|y^2|}{g(z)} = \frac{-\det B}{8}\frac{|f(z)|}{g^3(z)} \tag{9}$$

The total differential dK, is $dK(z) = \frac{\det B}{8}\frac{(-(g^3(z))d|f(z)|+3g^2|f|dg(z))}{(g^6(z))}$, and the differential dg is $dg(z) = b_{12} + b_{21} + b_{22}(d|(z - z_0)|^2)$. For $z = z_0$, we have that $dg(z)|_{(z=z_0)} = 0$ if and only if $b_{12} + b_{21} = 2\text{Re } b_{12} = 0$, then

$$dK(z)|_{(z=z_0)} = \frac{\det B}{8}\frac{(-g^3(z_0)(d|f(z)| + 3g^2(z_0)|f(z_0)|dg(z)|)}{(g^6(z_0))}|_{(z=z_0)}.$$

By hyphotesis $f'(z_0) = 0$, then $dK(z_0) = 0$ if and only if Re $b_{12} = 0$. Since this computation is local, we have that $(z_0, y_0) \in C$ is a critical point of K if and only if Re $b_{12} = 0$.■

Note that is easy to see that with the same hyphotesis the lemma 2.1 can be generalized for any hyperelliptic Riemann surface of genus $g > 0$.

Remark 2. The condition $b_{12} = \overline{b_{21}}$ and $b_{12} + b_{21} = 0 = \text{Re } b_{12}$ implies that $0 = (b_{12} + b_{21})^2 = b_{12}^2 + 2b_{12}b_{21} + b_{21}^2$, that is, $-4b_{12}b_{21} = 2(b_{12}^2 + b_{21}^2)$. Note that $i\text{Im } b_{12} = -i\text{Im } b_{21}$, so $b_{12}^2 + b_{21}^2 = ((i\text{Im } b_{12})^2 + (i\text{Im } b_{21})^2) = -((\text{Im } b_{12})^2 + (\text{Im } b_{21})^2) = (i\text{Im } b_{12} - i\text{Im } b_{21})(i\text{Im } b_{12} + i\text{Im } b_{21}) = 0$, that is,$-4b_{12}b_{21} = 0$, then $b_{12} = b_{21} = 0$, so $\det B = b_{11}b_{22} > 0$.

Lemma 2.2. With the same hypothesis of lemma 2.1, suppose also that $f''(z_0) = 0$, then (z_0, y_0) is a local minima of K of Morse index zero.

Proof. Let $\alpha := \frac{\det B}{8}$ be. From equation (9) we have that $\frac{\partial K}{\partial z} = -\alpha \cdot \frac{[g^3(z)\frac{\partial |f|}{\partial z} - |f(z)|3g^2(z)\frac{\partial g}{\partial z}]}{g^6(z)}$ and $\frac{\partial K}{\partial \overline{z}} = -\alpha \cdot \frac{[g^3(z)\frac{\partial |f|}{\partial \overline{z}} - |f(z)|3g^2(z)\frac{\partial g}{\partial \overline{z}}]}{g^6(z)}$. Then we have that

$$\frac{\partial^2 K}{\partial z^2} = -\alpha g^6(z)\frac{[3\frac{\partial|f|}{\partial z}g^2(z)\frac{\partial g}{\partial z} + g^3(z)\frac{\partial^2|f|}{\partial z^2} - 6|f(z)|g(z)(\frac{\partial g}{\partial z})^2 - 3g^2(z)(|f(z)|\frac{\partial^2 g}{\partial z^2} + \frac{\partial g}{\partial z}\frac{\partial|f|}{\partial z})]}{g^{12}(z)}$$
$$+ \alpha\frac{6g^5(z)\frac{\partial g}{\partial z}}{g^{12}(z)} \cdot [g^3(z)\frac{\partial|f|}{\partial z} - |f(z)|3g^2(z)\frac{\partial g}{\partial z}]$$
$$\frac{\partial^2 K}{\partial \overline{z}^2} = -\alpha g^6(z)\frac{[3\frac{\partial|f|}{\partial \overline{z}}g^2(z)\frac{\partial g}{\partial \overline{z}} + g^3(z)\frac{\partial^2|f|}{\partial \overline{z}^2} - 6|f(z)|g(z)(\frac{\partial g}{\partial \overline{z}})^2 - 3g^2(z)(|f(z)|\frac{\partial^2 g}{\partial \overline{z}^2} + \frac{\partial g}{\partial \overline{z}}\frac{\partial|f|}{\partial \overline{z}})]}{g^{12}(z)}$$
$$+ \alpha\frac{6g^5(z)\frac{\partial g}{\partial \overline{z}}}{g^{12}(z)} \cdot [g^3(s)\frac{\partial|f|}{\partial \overline{z}} - |f(z)|3g^2(z)\frac{\partial g}{\partial \overline{z}}]$$
$$\frac{\partial^2 K}{\partial \overline{z}\partial z} = \alpha\frac{[6g^5(z)\frac{\partial g}{\partial \overline{z}} \cdot [g^3(z)\frac{\partial|f|}{\partial z} - |f(z)|3g^2(z)\frac{\partial g}{\partial z}]}{g^{12}(z)} - \alpha g^6(z)\frac{[3g^2(z)\frac{\partial|f|}{\partial z}\frac{\partial g}{\partial \overline{z}} + g^3(z)\frac{\partial^2|f|}{\partial \overline{z}\partial z}]}{g^{12}(z)}$$
$$- \alpha g^6(z)[\frac{-|f(z)|(6g(z)\frac{\partial g}{\partial \overline{z}} \cdot \frac{\partial g}{\partial z} + 3g^2(z)\frac{\partial^2 g}{\partial \overline{z}\partial z}) - 3g^2(z)\frac{\partial g}{\partial z} \cdot \frac{\partial|f|}{\partial \overline{z}}]}{g^{12}(z)}$$

Note that $\frac{\partial^2|f|^2}{\partial z^2} = 2(\frac{\partial|f|}{\partial z})^2 + 2|f|\frac{\partial^2|f|}{\partial z^2}$. By hypthotesis $f(z_0) \neq 0, f'(z_0) = f''(z_0) = 0$, then $\frac{\partial^2|f|^2}{\partial z^2}(z_0) = 0$ and $\frac{\partial|f|}{\partial z}(z_0) = 0 = \frac{\partial|f|}{\partial \overline{z}}(z_0)$. Is easy to see that $\frac{\partial^2|f|}{\partial z^2}(z_0) = 0 = \frac{\partial^2|f|}{\partial \overline{z}\partial z}(z_0)$. Since $\frac{\partial^2 g}{\partial z^2}(z_0) = 0 = \frac{\partial^2 g}{\partial \overline{z}^2}(z_0)$ and $g(z_0) = b_{11}, \frac{\partial g}{\partial z}(z_0) = b_{21} = 0, b_{12} = \frac{\partial g}{\partial \overline{z}}(z_0) = b_{12} = 0$, we have from Remark 2:

$$\frac{\partial^2 K}{\partial z^2}(z_0) = \frac{[6\alpha|f(z_0)|b_{11}b_{21}^2]}{g^6(z_0)} - \frac{[18\alpha|f(z_0)|b_{11}b_{21}^2]}{g^6(z_0)} = \frac{-12\alpha|f(z_0)|b_{11}b_{21}^2}{g^6(z_0)} = 0.$$
$$\frac{\partial^2 K}{\partial \overline{z}^2}(z_0) = \frac{[6\alpha|f(z_0)|b_{11}b_{12}^2]}{g^6(z_0)} - \frac{[18\alpha|f(z_0)|b_{11}b_{12}^2]}{g^6(z_0)} = \frac{-12\alpha|f(z_0)|b_{11}b_{12}^2}{g^6(z_0)} = 0.$$
$$\frac{\partial^2 K}{\partial \overline{z}\partial z}(z_0) = \frac{[-18\alpha|f(z_0)|b_{12}b_{21}]}{g^5(z_0)} + \frac{6\alpha|f(z_0)|b_{12}b_{21}}{g^5(z_0)} + \frac{3\alpha|f(z_0)|b_{11}b_{22}}{g^5(z_0)}$$
$$= \frac{3\alpha|f(z_0)|}{g^5(z_0)}b_{11}b_{22} > 0.$$

Let $\delta := 2\frac{\partial^2 K}{\partial \overline{z}\partial z}(z_0)$ be. From (8), we have that the real Hessian of K in (z_0, y_0) is

$$\text{Hess } K(z_0, y_0) = \begin{pmatrix} \delta & 0 \\ 0 & \delta \end{pmatrix}.$$

This show that (z_0, y_0) is a non-degenerated critical point of Morse index 0, which is a local minima. ■

Example 3. Consider the hyperelliptic Riemann surface C of genus $g = 2$ defined by the equation $y^2 = z^6 - 1$. Let $\pi : C \to \mathbb{P}^1$ be the double cover ramified at Weierstrass points of C and consider the automorphism $\varphi : C \to C$,$\varphi(z, y) = (\zeta z, y)$, where $\zeta = e^{\frac{2\pi i}{6}}$. We are going to see that the curvature function $K : C \to \mathbb{R}$

of the Theta metric is a Morse function. To show this we we need a straightforward computation of partial derivates for K.

Note that the points $\{q_1 = (0, i), q_2 = (0, -i)\} = \pi^{-1}(0)($ the points lying over $0 \in \mathbb{P}^1)$ are fixed points of φ, so from lemma 1.1 they are critical points of K. From lemma 1.1 the automorphism $f : C \to C, f(z, y) = (\frac{1}{z}, \frac{iy}{z^3})$ leaves invariant K and from example two of section one K descends to $\mathbb{P}^1$ and $K(z) = K(\frac{1}{z})$. So we have also that the two points lying over $\infty \in \mathbb{P}^1$ are critical points of K. From lemma 2.1 and lemma 2.2 we have that $z_0 = 0$ is a local minima of K of Morse index zero, this implies that $\infty \in \mathbb{P}^1$ is a local minima of Morse index zero. Using the double cover π, we obtain four minimal points on C of Morse index zero. From Remark 2, $b_{12} = 0 = b_{21}$, so from equation (9), $K(z) = -\alpha \frac{|z^6-1|}{b_{22}^3(|z|^2+\frac{|b_{11}|^2}{|b_{22}|^2})^3}$. Since $K(z) = K(\frac{1}{z})$, a simple computation shows that $\frac{b_{11}}{b_{22}} = 1$. We obtain that $\alpha = \frac{\det\ B}{8} = \frac{b_{11}b_{22}}{8} = \frac{b_{22}^2}{8}$, that is, $K(z) = -\tau \cdot \frac{|z^6-1|}{(|z|^2+1)^3}$, where $\tau = \frac{1}{8b_{22}}$. Consider the function K^2. Since $d(K^2) = 2K \cdot dK$, except at the Weierstrass points, the differentials dK^2 and dK have the same critical points. In polar coordinates $z = re^{i\theta}$, $r > 0, \theta \in (0, 2\pi)$ we have that $K^2(z) = \tau^2 \frac{(z^6-1)\overline{(z^6-1)}}{(|z^2|+1)^6} = K^2(r, \theta) = \tau^2 \frac{r^{12}-2r^6 cos(6\theta)+1}{(r^2+1)^6}$. With respect to r and θ consider the partial derivates:

$$\begin{aligned}
\frac{1}{\tau^2} \cdot \frac{\partial K^2(r,\theta)}{\partial r} &= \frac{12r^{11} - 12r^5 cos(6\theta)}{(r^2+1)^6} - \frac{12r(r^{12} - 2r^6 cos(6\theta) + 1)}{(r^2+1)^7} \\
\frac{1}{\tau^2} \cdot \frac{\partial K^2(r,\theta)}{\partial \theta} &= \frac{12r^6 sin(6\theta)}{(r^2+1)^6}
\end{aligned}$$

All critical points of dK^2 are the point 0, the roots of $z^6 - 1 = 0$ and the points $(1, \theta_k)$ where $\theta_k = \frac{(2k+1)\pi}{6}$, $k = 0, 1, \ldots, 5$. The points $(1, \theta_k)$ corresponds to the complex number $\beta_k = \zeta^k \beta_1$, where $\zeta = e^{\frac{i\pi}{3}}, \beta_1 = e^{\frac{i\pi}{6}}, k = 0, 1, \ldots, 5$. The points β_k are the roots of $z^6 + 1 = 0$. A straightforward computation gives

$$\begin{aligned}
\frac{1}{\tau^2}\frac{\partial K^2}{\partial z} &= \frac{(|z^2|+1)^6 6z^5\overline{(z^6-1)} - 12\overline{z}(|z|^2+1)^5(z^6-1)\overline{(z^6-1)}}{(|z^2|+1)^{12}} \\
\frac{1}{\tau^2} \cdot \frac{\partial^2 K^2}{\partial^2 z^2} &= \frac{30z^4(\overline{z^6-1})}{(|z|^2+1)^6} - \frac{144|z^2|z^4(\overline{z^6-1})}{(|z|^2+1)^7} + \frac{168\overline{z}^2|z^6-1|^2}{(|z|^2+1)^8} \\
&= \frac{1}{\tau^2} \cdot \frac{\partial^2 K^2}{\partial^2 \overline{z}^2} \\
\frac{1}{\tau^2} \cdot \frac{\partial^2 K^2}{\partial \overline{z} \partial z} &= \frac{36|z|^{10}}{(|z|^2+1)^6} + \frac{72(\overline{z}^6 - z^6) - 12|z^6-1|^2 - 144z^6(\overline{z^6-1})}{(|z|^2+1)^7} + \frac{168|z|^2|z^6-1|^2}{(|z|^2+1)^8}.
\end{aligned}$$

Since $\beta_1^6=-1$ then $\beta_1^4=\frac{-1}{2}+i\frac{\sqrt{3}}{2}$, $\beta_1^2=\frac{1}{2}+i\frac{\sqrt{3}}{2}$ and $\beta_1^4+\overline{\beta}_1^2=0$. This implies that

$$\frac{\partial^2K^2}{\partial z^2}(\beta_1)=\frac{84(\beta_1^4+\overline{\beta}_1^2)+84\overline{\beta}_1^2}{2^6}=\frac{84\overline{\beta}_1^2}{2^6},\ \frac{\partial^2K^2}{\partial\overline{z}\partial z}(\beta_1)=\frac{36}{2^6}.$$

Let $(u_0,v_0)=(\mathrm{Re}\ \beta_1,\mathrm{Im}\ \beta_1)$ be. From (8) we have in real coordinates that

$$\frac{\partial^2K^2}{\partial u^2}(u_0,v_0)=\frac{36+42(\overline{\beta}_1^2+\beta_1^2)}{2^5}\ ,\frac{\partial^2K^2}{\partial^2 v^2}(u_0,v_0)=\frac{36-42(\overline{\beta}_1^2+\beta_1^2)}{2^5},\ \frac{\partial^2K^2}{\partial u\partial v}(u_0,v_0)=\frac{42\sqrt{3}}{2^5},$$

Note that $\overline{\beta}_1^2+\beta_1^2=1,\beta_1^2-\overline{\beta}_1^2=i\sqrt{3}$, so

$$\mathrm{Hess}\ K(u_0,v_0)=\frac{\tau^2}{2^5}\begin{pmatrix}78 & 42\sqrt{3}\\ 42\sqrt{3} & -6\end{pmatrix}.$$

This implies that β_1 is a saddle point of Morse index $+1$. Note that $\varphi(\beta_j,y)=(\beta_{j+1},y)$. So each point β_j has Morse index $+1$. The roots of $z^6+1=0$ are not ramification points of the double cover $\pi:C\to\mathbb{P}^1$, then we have that the 12 points $\pi^{-1}(\{z^6+1=0\})$ are saddle points K of Morse index $+1$. Thus for the curvature function $K:C\to\mathbb{R}$, we have $I_0=4$ critical points of Morse index zero, $I_1=12$ critical points of Morse index $+1$ and $I_2=6$ critical points of Morse index $+2$. The Euler characteristic is $\mathcal{X}(C)=2-2g=-2=I_0-I_1+I_2$. This implies that K is a Morse function on C.■

REFERENCES

[1] P. Griffiths, J. Harris,*Principles of Algebraic Geometry* , Jhon Wiley and Sons, 1994.

[2] H. Lange, C. Birkenhake,*Complex Abelian varieties*, Second edition, Springer Verlag, 2004.

[3] J. Lewittes,*Differentials and metrics on Riemann surfaces*, Trans. Amer. Math. Soc. 139, 1969(311-318).

Abel Castorena
Instituto de Matemáticas, Unidad Morelia.
Universidad Nacional Autonoma de Mexico.
Apdo. Postal 61-3(Xangari). C.P. 58089.
Morelia, Michoacán. MEXICO.
e-mail address: abel@matmor.unam.mx